

# orphananesthesia

## Anaesthesia recommendations for patients suffering from **Lenz-Majewski hyperostotic dwarfism**

**Disease name:** Lenz-Majewski hyperostotic dwarfism

**ICD 10:** Q 87.1

**Synonyms:** Dystrophinopathy

Lenz-Majewski hyperostotic dwarfism is a very rare syndrome caused by a de novo heterozygous mutation of phosphatidylserine synthase 1 (PTDSS1) gene (Sousa et al. 2013). It was first described by Braham in 1969 as the Camurati-Engelman syndrome but soon later became the Braham-Lenz syndrome until it was re-classified by Lenz and Majewski in 1974. Only thirteen reported cases are described in the Anglo-Saxon literature.

The most important feature is a disproportionately large head in relation to a reduced size trunk and limbs. Large fontanels and widely separated sutures that close lately in childhood are present. The skin is loose, wrinkled, and atrophic with prominent veins, especially in the scalp. The ears are large and floppy, and frequently associated are choanal atresia or stenosis, nasolacrimal duct obstruction, and, in boys, cryptorchidism and inguinal hernia. There is craniofacial dysmorphic evolution with age, leading to progressive macroglossia, prognathism, midface hypoplasia, and upper airway obstruction.

---

Medicine in progress



Perhaps new knowledge

Every patient is unique

Perhaps the diagnostic is wrong

---

---

### Disease summary

---

The disorder is also characterized by failure to thrive and severe mental retardation. This condition can be mistaken for craniometaphyseal and craniodiaphyseal dysplasias. However, the absence of cranial nerves impingement confirms the diagnosis. Enlarged ventricles and hydrocephalus have been described in some patients.

The skeletal alterations are striking with the following radiographic features:

- progressive sclerosis of the skull, facial bones, and vertebrae
- broad clavicles and ribs
- short or absent middle phalanges
- diaphyseal undermodeling and midshaft cortical thickening
- metaphyseal and epiphyseal hypostosis
- retarded skeletal maturation

Tooth enamel is also defective.

---

### Typical surgery

---

Musculoskeletal disorders procedures are common.

---

### Type of anaesthesia

---

There are no specific precautions before anaesthesia except those associated with other medical conditions with the patient. However, because of the nature of the disease, facial dysmorphism is very common, and one must carefully evaluate the airway and plan accordingly for a potential difficult airway management and tracheal intubation.

This disease can involve considerable musculoskeletal and skin features that will require special attention during positioning. Finally, due to mental retardation, the use of a regional anaesthetic technique should be complemented by controlled sedation or general anaesthesia. In both cases, proper monitoring of the airway is mandatory.

No specific anaesthetic considerations are reported for this medical condition besides the usual patient management. However, in presence of a difficult airway, induction of anaesthesia with a volatile anaesthetic technique, maintenance of spontaneous breathing and placement of an intravenous catheter must be performed safely. Tracheal intubation under safe conditions is highly recommended. The use of a laryngeal mask is an alternative only for procedure where access to the airway is always available.

There are no known pharmacological implications for this syndrome.

Neuroaxial or peripheral block is an interesting option for anaesthetic management. The use of ultrasound-guided nerve blocks enhances success rate.

---

**Necessary additional diagnostic procedures (preoperative)**

---

Due to the low incidence of this syndrome and the lack of clinical evidence, no specific recommendation besides the usual preoperative preparation must be done. However, as the child gets older and thoracic deformity worsens, lung function tests might be considered (when feasible) to establish the pulmonary involvement.

---

**Particular preparation for airway management**

---

Not reported.

---

**Particular preparation for transfusion or administration of blood products**

---

Not reported.

---

**Particular preparation for anticoagulation**

---

Not reported.

---

**Particular precautions for positioning, transport or mobilisation**

---

Positioning might require special attention particularly because of leg problems and the exquisite pain manifested by these patients in the postoperative period.

---

**Probable interaction between anaesthetic agents and patient's long term medication**

---

Not reported.

---

**Anaesthesiologic procedure**

---

There are no pharmacological known implications with this medical condition.

---

**Particular or additional monitoring**

---

Standard monitoring is indicated. The use of invasive monitoring should be dictated by the surgical procedure alone and not the medical condition.

---

#### Possible complications

---

Although they are not directly reported, given facial dysmorphism, potential complications could be related to difficult airway management.

Postoperative exquisite pain has been reported postoperatively and should be an important consideration in the management of these patients.

---

#### Postoperative care

---

The need of postoperative care is related to the kind of surgery and to preexisting comorbidity.

---

#### Information about emergency-like situations / Differential diagnostics

---

*caused by the illness to give a tool to distinguish between a side effect of the anaesthetic procedure and a manifestation of the diseases, e.g.:*

Not reported.

---

#### Ambulatory anaesthesia

---

There are no contraindications to same-day anaesthesia and surgical procedures. Each patient must be evaluated carefully for co-morbidity and/or airway issues that might suggest that a hospitalization will be necessary for the safety of the postoperative care.

---

#### Obstetrical anaesthesia

---

Not reported.

### Literature and internet links

1. Sousa SB, Jenkins D, Chanudet E, Tasseva G, Ishida M, Anderson G, Docker J, Ryten M, Sa J, Saraiva JM, Barnicoat A, Scott R, Calder A, Wattanasirichaigoon D, Chrzanowska K, Simandlova M, Van Maldergem L, Stanier P, Beales PL, Vance JE, Moore GE. Gain-of-function mutations in the phosphatidylserine synthase 1 (PTDSS1) gene cause Lenz-Majewski syndrome. *Nat Genet.* 2014 Jan;46(1):70-6. doi: 10.1038/ng.2829. Epub 2013 Nov 17
2. Wattanasirichaigoon D, Visudtibhan A, Jaovisidha S, Laothamatas J, Chunharas A. Expanding the phenotypic spectrum of Lenz-Majewski syndrome: facial palsy, cleft palate and hydrocephalus. *Clin Dysmorphol.* 2004 Jul;13(3):137-42
3. Gorlin RJ, Whitley CB. Lenz-Majewski syndrome. *Radiology.* 1983 Oct;149(1):129-31
4. Robinow M, Johanson AJ, Smith TH. The Lenz-Majewski hyperostotic dwarfism. A syndrome of multiple congenital anomalies, mental retardation, and progressive skeletal sclerosis. *J Pediatr.* 1977 Sep;91(3):417-21
5. Lenz WD, Majewski F. A generalized disorders of the connective tissues with progeria, choanal atresia, symphalangism, hypoplasia of dentine and craniodiaphyseal hypostosis. *Birth Defects Orig Artic Ser.* 1974;10(12):133-6
6. Baum VC, O'Flaherty J. *Anesthesia for Genetic, Metabolic, and Dysmorphic Syndromes of Childhood.* 2<sup>nd</sup> edition, Philadelphia: Lippincott Williams & Wilkins; 2006
7. Bissonnette B, Luginbuehl I, Marciniak B, Dalens B. *Syndromes – Rapid recognition and perioperative implications.* McGraw-Hill, New York; 2007.

---

**Last date of modification: August 2014**

---

*These guidelines have been prepared by:*

**Authors**

**Adrià Font Gual**, Anaesthesiologist, Hospital de la Santa Creu i Sant Pau. Barcelona, Spain  
[AFontG@santpau.cat](mailto:AFontG@santpau.cat)

**Tobias Koller**, Anaesthesiologist, Hospital de la Santa Creu i Sant Pau. Barcelona, Spain

**Juan Ignacio Casas Vila**, Head of Unit, Department of Anaesthesiology. Hospital de la Santa Creu i Sant Pau. Barcelona, Spain

**Peer revision 1**

**Bruno Bissonnette**, Director of Neurosurgical Anesthesia, University of Toronto, Canada  
[bruno@brainstorm-anesthesia.com](mailto:bruno@brainstorm-anesthesia.com)

**Peer revision 2**

**Duang Wattanasirichaigoon**, Paediatrician, Ramathibodi Hospital, Mahidol University, Bangkok, Thailand  
[duangrurdee.wat@mahidol.ac.th](mailto:duangrurdee.wat@mahidol.ac.th)

---