

# orphananesthesia

## Anaesthesia recommendations for patients suffering from Myasthenia gravis

**Disease name:** Myasthenia gravis (MG)

**ICD 10:** G70.0

**Synonyms:** Myasthenia gravis (juvenile and adult form) autoimmune (receptor-binding antibodies)

Additional forms of myasthenia:

- Isolated ocular myasthenia (autoimmune)
- Congenital myasthenia (presynaptic, postsynaptic)
- Neonatal myasthenia (maternal antibodies in newborns)

Myasthenia gravis (MG) is an autoimmune disorder with increasing frequency and recognition and is present in the paediatric and adult population. The juvenile form of MG is the most common and is similar to the adult form. It is caused by antibodies against the acetylcholine receptor at the postsynaptic neuromuscular junction. 75% of these cases occur after the age of ten [1]. The clinical expression differentiates between the ocular and the generalized form. The first clinical signs are painless weakness followed by ptosis and diplopia. Ptosis and diplopia images as the only clinical presentation occur in the isolated ocular form of myasthenia gravis. If muscular weakness progresses within the next year, a generalized form is present with the potential for respiratory muscle involvement. The first line of treatment are oral anticholinesterase inhibitors, such as pyridostigmine. However, high doses could induce cholinergic crises. Thymectomy mainly as an endoscopic procedure is currently done in young onset acetylcholine-receptor antibody positive patients with generalized myasthenia [2,3].

Due to the immunological origin of the disorder, immune suppressant substances could be helpful. Mainly steroids are used, but also azathioprine, ciclosporin, methotrexate and cyclophosphamide could be used in poor-responders or as steroid-sparing agents for chronic use.

---

Medicine in progress



Perhaps new knowledge

Every patient is unique

Perhaps the diagnostic is wrong

---

### Disease summary

Congenital myasthenic syndromes are genetically, mainly recessive, transmitted disorders. This group is clinically very heterogeneous, children present shortly after birth with a feeding problem and muscular hypotonia. As a severe symptom, respiratory insufficiency is well described, and children often need artificial ventilation. A few children respond to therapy with acetylcholinesterase-blockers. The defect could be located presynaptic (e.g. Cholinacetyl-Transferase Defect CHAT), synaptic (e.g. mutations in the gene encoding for Collagen Q COLQ) and postsynaptic (e.g. disturbances in fast or slow channels in acetylcholinrezeptors or mutations in the gene encoding for the Rapsyn gene). A special feature in CHAT defects are recurrent apnea in children suffering from simple infectious diseases. These children require fast intubation and artificial ventilation.

Transient neonatal myasthenia gravis occurs in approximately 12% of mothers with MG and improves within the first weeks of live as the antibody concentration decreases. The first signs are often muscular hypotonia, feeding problems and respiratory distress.

The characteristic presentation of MG is fatigability of voluntary muscles – depending on the extent of the disease –, which typically worsens with continued daily activity.

### Typical surgery

- Thymectomy
- General surgery
- Obstetrics and gynecology
- Tracheostomy

### Type of anaesthesia

Locoregional anaesthesia is possible, depending on surgery. There is no evidence regarding the role of local anaesthetics, but theoretically, ester local anaesthetics might lead to problems in patients with MG and an anticholinesterase medication due to inhibition of metabolism [4]. High blood levels of any of the local anaesthetics can cause muscular weakness.

Epidural techniques with bupivacaine in patients with MG and transsternal thymectomy had been reported uneventful. Spinal anaesthesia is possible [5].

Analgo-sedation should be performed very carefully as any sedative or muscle relaxing medication can cause respiratory depression (Table 1).

General anaesthesia is possible and necessary.

Induction with propofol, barbiturates, etomidate or ketamine are described as uneventful in patients with MG [6,7]. Opioids analgesics do not depress neuromuscular transmission, and remifentanyl's short elimination half-time makes it more titratable in myasthenic patients [8].

Benzodiazepines should be avoided due to increase of myasthenic symptoms.

Volatile anaesthetics can be used but may cause muscle relaxation and show reduced train of four (TOF) responses in MG patients like in all other patients. There is no difference in the use of sevoflurane or isoflurane [12,13].

The myasthenic patient is very sensitive to any neuromuscular blockade. The use of small doses for priming or defasciculation is not appropriate as the use of long acting muscle relaxants like pancuronium. In myasthenic patients the ED95 of vecuronium has an individual variability and varies between 5% to 90% of the normal dose; the same with atracurium and cisatracurium.

There is an increased sensitivity to any non-depolarizing muscle relaxants. Dose has to be reduced by 2 to 8 times. The TOF ratio before the administration of the muscle relaxant can give an adequate information of the needed dose, but still the duration of the neuromuscular can be prolonged.

Succinylcholine as a depolarizing muscle-relaxing agent can be used, but be aware that patients with MG have a reduced response and need higher doses of **succinylcholine** 1.5 to 2.0 mg/kg.

The metabolism of mivacurium and succinylcholine can be affected by the preoperative use of pyridostigmine, and therefore the duration of the neuromuscular blockade can be prolonged [9,10,11].

A good alternative might be the intubation with propofol and remifentanyl.

Antagonization of muscle relaxants may be difficult due to risk of cholinergic crisis and must be titrated carefully.

#### **Necessary additional diagnostic procedures (preoperative)**

##### Basic medication

- Anticholinesterase medication could be stopped, if this does not influence the respiratory situation. The need for muscle relaxants during the operation can be reduced
- Steroids: Should be given and covered preoperatively
- Immuno suppressant

##### Blood count and electrolytes

Lung function test should be done preoperatively (classification of MG [18]) and blood gas analysis. Chest X-Ray or CT-Scan before thymectomy for detection of tracheal compression or deviated trachea.

Plasmapheresis could be useful in patients with autoimmune triggered MG (juvenile or adult form). This is typically performed prior to surgery in patients with unstable disease.

Premedication should be avoided in patients with severe reduced respiratory reserve, in cases with isolated ocular form of MG reduced doses of benzodiazepine are acceptable.

### Particular preparation for airway management

---

No specific problems expected. Be aware of the possibility of postoperative need of mechanical ventilation or prolonged time of extubation.

### Anaesthesiologic procedure

---

If possible, regional techniques should be preferred. General anaesthesia as balanced or total intravenous anaesthesia can be used if reduced doses of muscle relaxants are used.

In case of thymectomy, epidural techniques for intra- and postoperative analgesia should be intended.

### Particular or additional monitoring

---

Standard monitoring for normal operative procedure and extended monitoring according to the underlying operation.

Monitoring of the neuromuscular blockade is mandatory:

The preferred monitoring of neuromuscular transmission is electromyogram (EMG) and mechanomyogram (MMG) or calibrated acceleromyography (AMG) and train of four (TOF). TOF should be performed in the awake patient preoperatively, and the ratio of T4/T1 (fade ratio) should be 1.0. In MG patients the ratio could be < 0.9 and shows a reduced need for depolarizing muscle relaxants. The use of calibrated AMG is recommended, and the TOF ratio should reach preoperative values [15].

### Postoperative care

---

Should be adapted to the underlying operations and comorbidity. The need for postoperative ventilation is dependent on the operation and the preoperative clinical extent of the MG. Possible criteria are a duration disease longer than 6 years, respiratory complications in the past medical history and a reduced vital capacity < 50% according to age.

Adequate postoperative pain and pain control, pulmonary toilet and the avoidance of drugs that interfere with neuromuscular transmission (Table 1). Anticholinesterase medication has to be continued in the postoperative phase. The dose has to be titrated, and the intravenous dose if needed has to be in an equivalent dosage.

Patients should stay in a postanesthesia care unit or an intensive care unit.

Lists, summarizing substances which are definitely deteriorating myasthenia gravis are available (table 1, [www.dgn.org](http://www.dgn.org), [www.myasthenia.org](http://www.myasthenia.org)). Specially antibiotics, anticonvulsant drugs, beta-blocker and psychopharmacological medications are of special interest.

---

#### **Ambulatory anaesthesia**

---

Unsafe and should be avoided.

---

#### **Obstetrical anaesthesia**

---

Obstetrical anaesthesia can be performed as general as well as regional anaesthesia. Like in most neuromuscular disorders, MG can deteriorate during pregnancy [16].

Maternal antibodies in MG can pass the placenta barrier and give the possibility for impaired newborns with muscular hypotonia or respiratory distress after birth.

## Annex 1

Table 1. Drugs with possible influence on myasthenia gravis

Class	Drugs
Analgesics	flupirtine, morphine
Antiarrhythmics	chinidine, ajmaline, procainamide, mexitile, $\beta$ -blocker (pindolol, propranolol, timolol topical)
Antibiotics	aminoglycosides, macrolide, ketolide, lincomycine, fluoroquinolone, sulfonamide, tetracycline
Antidepressants	amtryptiline, lithium
Anticonvulsants	benzodiazepine, carbamazepine, ethosuximide, gabapentin
Antimalarial Drugs	chinidine, chloroquine
Antirheumatic Drugs	d-penicillamine, chloroquine
Ca-channel Inhibitor	verapamil, nifedipine, diltiazem
Diuretics	loop-diuretics, acetazolamide
Local anaesthetics	ester type
Magnesium	high dose therapy (e.g. obstetrics)
Psychopharmaceutical	chlorpromazine and other phenothiazines, benzodiazepine

### Literature and internet links

1. Fenichel GM. Myasthenia gravis. *Pediatr Anaesth* 1989;18(7):432-8
2. Adams C, Murphy E. Thymectomy in juvenile myasthenia gravis. *J Child Neurol* 1990;5:215-218
3. Rodriguez M, Howard F. Myasthenia gravis in children: Long-term follow up. *Ann Neurol* 1983; 13:504-510
4. Matsuo S, et al. Interaction of muscle relaxants and local anesthetics at the neuromuscular junction. *Anesth Analg* 1978;57(5):580-7
5. Kawamata M, et al. Continuous thoracic epidural blockade in combination with general anesthesia with nitrous oxide, oxygen, and sevoflurane in two patients with myasthenia gravis. *Masui* 1993;42(6):898-901
6. O'Flaherty D, et al. Total intravenous anesthesia with propofol for transsternal thymectomy in myasthenia gravis. *J Clin Anesth* 1992;4(3):241-4
7. Florence AM. Anaesthesia for transcervical thymectomy in myasthenia gravis. *Ann R Coll Surg Engl* 1984;66(5):309-12
8. Fodale V, et al. Propofol and remifentanyl without muscle relaxants in a patient with myasthenia gravis for emergency surgery. *Can J Anaesth* 2003;50(10):1083-4
9. Nilsson E, Meretoja O. Vecuronium dose-response and maintenance requirements in patients with myasthenia gravis. *Anesthesiology* 1990;73(1):28-32
10. Baraka A. Anesthesia and critical care of thymectomy for myasthenia gravis. *Chest Surg Clin N Am* 2001;11(2):337-61
11. Eisenkraft JB, Papatestas AE. Anesthesia for trans-sternal thymectomy in myasthenia gravis. *Ann R Coll Surg Engl* 1988;70(4):257-8
12. Kadosaki M, et al. Severity of myasthenia gravis is related to the degree of neuromuscular blocking effect by isoflurane. *Masui* 1993;42(6):906-9
13. Itoh H, Shibata K. Comparison between sevoflurane and propofol neuromuscular effects in a patient with myasthenia gravis: Effective doses of vecuronium. *Anesthesiology* 2001;95(3): 803-5
14. Osserman KE, Jenkins G. Studies in myasthenia gravis: Review of a twenty-year experience in over 1200 patients. *Mt Sinai J Med* 1971;38(6):497-537
15. Murphy GS, et al. Intraoperative acceleromyography reduces symptoms of muscle weakness and improves quality of recovery in the early postoperative period. *Anesthesiology* 2011; 115(5):946-954
16. Sax TW, Rosenbaum RB. Neuromuscular disorders in pregnancy. *Muscle Nerve* 2006;12
17. Mann R, Blobner M, Jelen-Esselborn S, Busley R, Wener C. Preanesthetic train-of-four fade predicts the atracurium requirement of myasthenia gravis patients. *Anesthesiology* 2000; 93(2):346-350
18. Besinger UA, Toyka KV. Myasthenia gravis: Long-term correlation of binding and bungatoxin blocking antibodies against acetylcholin receptors with changes in disease severity. *Neurology* 1983;33:1316-1321.

---

**Last date of modification: August 2014**

---

*These guidelines have been prepared by:*

**Author**

**Thomas Fischer**, Anaesthesiologist, Zentrum für Frauen- und Kindermedizin Kassel,  
Germany  
[t.fischer@klinikum-kassel.de](mailto:t.fischer@klinikum-kassel.de)

**Peer revision 1**

**Tino Muenster**, Anaesthesiologist, University Hospital Erlangen, Germany  
[tino.muenster@kfa.imed.uni-erlangen.de](mailto:tino.muenster@kfa.imed.uni-erlangen.de)

**Peer revision 2**

**Nicholas Silvestri**, Neurologist, Adult Neurology Residency University, Buffalo School of  
Medicine and Biomedical Sciences, Buffalo, USA  
[njs6@buffalo.edu](mailto:njs6@buffalo.edu)

---