

A&I

ANÄSTHESIOLOGIE & INTENSIVMEDIZIN

Offizielles Organ: Deutsche Gesellschaft für Anästhesiologie und Intensivmedizin e.V. (DGAI)
Berufsverband Deutscher Anästhesisten e.V. (BDA)
Deutsche Akademie für Anästhesiologische Fortbildung e.V. (DAAF)
Organ: Deutsche Interdisziplinäre Vereinigung für Intensiv- und Notfallmedizin e.V. (DIVI)



Mitochondrial disorders
Neuronal ceroid lipofuscinosis

orphan**a**nesthesia

a project of the German Society
of Anaesthesiology and Intensive Care Medicine

SUPPLEMENT NR. 6 | 2017

OrphanAnesthesia –

ein krankheitsübergreifendes Projekt des Wissenschaftlichen Arbeitskreises Kinderanästhesie der Deutschen Gesellschaft für Anästhesiologie und Intensivmedizin e.V.

Ziel des Projektes ist die Veröffentlichung von Handlungsempfehlungen zur anästhesiologischen Betreuung von Patienten mit seltenen Erkrankungen. Damit will Orphan Anesthesia einen wichtigen Beitrag zur Erhöhung der Patientensicherheit leisten.

Patienten mit seltenen Erkrankungen benötigen für verschiedene diagnostische oder therapeutische Prozeduren eine anästhesiologische Betreuung, die mit einem erhöhten Risiko für anästhesieassoziierte Komplikationen einhergehen. Weil diese Erkrankungen selten auftreten, können Anästhesisten damit keine Erfahrungen gesammelt haben, so dass für die Planung der Narkose die Einholung weiterer Information unerlässlich ist. Durch vorhandene spezifische Informationen kann die Inzidenz von mit der Narkose assoziierten Komplikationen gesenkt werden. Zur Verfügung stehendes Wissen schafft Sicherheit im Prozess der Patientenversorgung.

Die Handlungsempfehlungen von OrphanAnesthesia sind standardisiert und durchlaufen nach ihrer Erstellung einen Peer-Review-Prozess, an dem ein Anästhesist sowie ein weiterer Krankheitsexperte (z.B. Pädiater oder Neurologe) beteiligt sind. Das Projekt ist international ausgerichtet, so dass die Handlungsempfehlungen grundsätzlich in englischer Sprache veröffentlicht werden.

Ab Heft 5/2014 werden im monatlichen Rhythmus je zwei Handlungsempfehlungen als Supplement der A&I unter www.ai-online.info veröffentlicht. Als Bestandteil der A&I sind die Handlungsempfehlungen damit auch zitierfähig. Sonderdrucke können gegen Entgelt bestellt werden.

OrphanAnesthesia –

a common project of the Scientific Working Group of Paediatric Anaesthesia of the German Society of Anaesthesiology and Intensive Care Medicine

The target of OrphanAnesthesia is the publication of anaesthesia recommendations for patients suffering from rare diseases in order to improve patients' safety. When it comes to the management of patients with rare diseases, there are only sparse evidence-based facts and even far less knowledge in the anaesthetic outcome. OrphanAnesthesia would like to merge this knowledge based on scientific publications and proven experience of specialists making it available for physicians worldwide free of charge.

All OrphanAnesthesia recommendations are standardized and need to pass a peer review process. They are being reviewed by at least one anaesthesiologist and another disease expert (e.g. paediatrician or neurologist) involved in the treatment of this group of patients.

The project OrphanAnesthesia is internationally oriented. Thus all recommendations will be published in English.

Starting with issue 5/2014, we'll publish the OrphanAnesthesia recommendations as a monthly supplement of A&I (Anästhesiologie & Intensivmedizin). Thus they can be accessed and downloaded via www.ai-online.info. As being part of the journal, the recommendations will be quotable. Reprints can be ordered for payment.

Bisher in A&I publizierte Handlungsempfehlungen finden Sie unter:

www.ai-online.info/Orphsuppl
www.orphananesthesia.eu

A survey of until now in A&I published guidelines can be found on:

www.ai-online.info/Orphsuppl
www.orphananesthesia.eu



Deutsche Gesellschaft für Anästhesiologie & Intensivmedizin

www.dgai.de



ANÄSTHESIOLOGIE & INTENSIVMEDIZIN

www.ai-online.info

Projektleitung

Prof. Dr. Tino Münster, MHBA
Geschäftsführender Oberarzt
Facharzt für Anästhesie,
Spezielle Schmerztherapie,
Notfallmedizin
Anästhesiologische Klinik
Friedrich-Alexander-Universität
Erlangen-Nürnberg
Krankenhausstraße 12
91054 Erlangen, Deutschland
Tel.: 09131 8542441
Fax: 09131 8536147
E-Mail: muenster@kfa.imed.uni-erlangen.de

orphananesthesia

Anaesthesia recommendations for patients suffering from **Neuronal ceroid lipofuscinosis**

Disease name: Neuronal ceroid lipofuscinoses (NCL)

ICD 10: E75.4

Synonyms: Historically, single NCL forms have been classified according to infantile, late-infantile, juvenile or adult onset and associated with names of investigators such as Santavuori-Haltia, Jansky-Bielschowsky, Batten, Spielmeyer-Vogt, Kufs.

The neuronal ceroid lipofuscinoses (NCL) are a heterogeneous group of genetic lysosomal storage diseases causing dementia, epilepsy, motor deterioration and mostly also visual loss through retinal degeneration. Similarities of the different NCL also concern elements of their pathophysiology, which is characterized by loss of neurons and accumulation of a material called ceroid lipofuscin. The single NCL forms differ significantly by the age at manifestation, and the progression of neurological deterioration.

Medicine in progress



Perhaps new knowledge

Every patient is unique

Perhaps the diagnostic is wrong

► **Citation:** Richa F: Neuronal ceroid lipofuscinosis. AnästH Intensivmed 2017;58:S134-S140. DOI: 10.19224/ai2017.S134

Disease summary

The neuronal ceroid lipofuscinoses (NCL) are a heterogeneous group of genetic lysosomal disorders characterized by the accumulation of a waxy intracellular storage material termed ceroid lipofuscin and progressive neurological deterioration, usually associated with dementia and epilepsy, frequently also with visual loss due to retinopathy. Most NCL forms are transmitted autosomal recessively. As a group, NCL represent one of the most frequent etiologies of dementia in children and young adults [1-3]. The classification of the NCL disorders is based on the different genes involved (CLN1-CLN14) and on the age at clinical onset, which can be anytime between the infantile and young adult age. The major NCL forms are shortly described below according to the age at manifestation.

NCL with infantile onset (1st year of life)

Infantile NCL is usually caused by CLN1 mutations. Early development appears normal until around six months of age. At onset, there is typically a decreased muscle tone and decreased social interaction followed by dramatically progressive psychomotor decay, myoclonus, seizures, spasticity and visual failure. By two years of age, blindness has developed with optic atrophy and retinal abnormalities. Patients rapidly reach a vegetative state.

NCL with late-infantile onset (age 2 to 5 years)

The onset of an NCL disease at this age is usually caused by CLN2 mutations, but mutations of other genes are possible. Children with CLN2 disease appear initially healthy and normally developed. Acquisition of speech may be retarded. First symptoms occur generally between the ages of 2 and 4 years and consist in motor decline with clumsiness and ataxia, deterioration of speech and/or epilepsy. Seizures (partial, generalized tonic-clonic, secondarily generalized, or other) are the first symptom in 75% of patients. Non-epileptic myoclonus frequently coexists and has to be distinguished from epileptic seizures as it is treated differently. After the third year of life, loss of motor function and of language ability progress rapidly and uniformly. MR imaging shows progressive brain atrophy. Visual ability declines gradually and leads to blindness. Limb spasticity, truncal hypotonia, and loss of head control lead to complete loss of independent mobility. Children lose the ability to swallow and frequently receive tube feeding. Death usually occurs at the age of 10-15 years.

NCL with juvenile onset (age 5 to 15 years)

Juvenile NCL is mostly caused by CLN3 mutations, rarely also by mutations of other genes. The disease usually starts between ages 4 and 7 years with insidious onset of visual failure due to a pigmentary retinopathy. For several years, patients are regarded as visually handicapped only. Progressive cognitive decline and behavioral problems (angry outbursts, physical violence, and anxiety with features of depression) follow. Seizures develop at around 10 years of age, mostly as generalized tonic-clonic seizures, which are usually well-controlled by medication, at least initially. Any other type of epilepsy may occur, from subtle partial seizures to myoclonic status. A parkinsonian movement disorder develops. Speech becomes affected by a severe dysarthria. Swallowing difficulties frequently lead to tube feeding. A cardiac conduction abnormality is detectable in the second decade of life. Age at death is usually in the third decade.

NCL with onset in adults

Adult onset forms of NCL are all very rare and can be caused by mutations of a variety of genes. In the past, the term Kufs disease was used to designate some of them. Onset is

typically at around 30 years of age. Clinical features include ataxia, dementia, progressive myoclonus epilepsy, but in contrast to most other NCL, there is usually no loss of vision.

Typical surgery

Children who have NCL often require anaesthetic care for diagnostic procedures, gastrostomy, vagus nerve stimulation or other types of surgery. There is limited information available concerning the anaesthetic management of a patient with NCL [4-6]. Because of the profound neurologic abnormalities in these children, the anaesthetic management of these patients requires an understanding of the natural history of the disease.

Type of anaesthesia

In the literature, there is insufficient data to recommend a specific type of anaesthesia. NCL patients are frequently blind, demented and tend to have seizures. They may have severe neurologic abnormalities, including difficulty swallowing with the consequence of an increased risk for aspiration. Patients are prone to bradycardia and may develop hypothermia during anaesthesia.

Necessary additional diagnostic procedures (preoperative)

Preoperative assessment should be conducted with special focus on epilepsy and other coexisting medical problems. The type and pattern of seizures, frequency and control of seizures, antiepileptic therapy, and complications of antiepileptic drugs (AEDs) need to be sought. Laboratory tests should screen for the side-effects of AEDs. Patients on sodium valproate, carbamazepine and ethosuximide should have their liver function, platelet count, and coagulation indices checked. In poorly controlled epilepsy, plasma levels of AEDs should be measured and optimized before operation. Patients on valproate may develop a severe exacerbation of a complex movement disorder leading to hyperthermia, hyper-CK-emia and prolonged decreased consciousness [7]. Electrocardiogram is necessary to rule out cardiac conduction disorders. X-ray or blood gas analysis should be considered in cases with suspected recurrent aspiration. Consultation with specialists may be advised (cardiology, paediatric neurology, respiratory care [8]).

Particular preparation for airway management

Preoperative physiotherapy may be advisable to remove excessive tracheobronchial secretions due to a swallowing disorder. There are no good data on the use of specific premedication.

Particular preparation for transfusion or administration of blood products

There are no particular recommendations.

Particular preparation for anticoagulation

The presence of an impaired mobility may predispose to deep venous thrombosis, so compression stockings and/or low molecular weight heparin may be indicated in the perioperative period.

Particular precautions for positioning, transport or mobilisation

These patients may suffer spinal rigidity, scoliosis and limb contractures, although uncommon; for this reason, positioning in the operating room must be careful. Immobile patients may have breakable bones, and fractures can be caused by inadequate handling.

Probable interaction between anaesthetic agents and patient's long-term medication

The use of preoperative AEDs may have significant impact on anaesthetic management. Cytochrome p450 enzyme induction by AEDs (e.g. phenytoin, a drug not recommended in NCL patients) may alter the metabolism of anaesthetic drugs. AEDs may also affect neuromuscular relaxants by causing acetylcholine receptors to be up-regulated at the neuromuscular junction.

Anaesthesiologic procedure

Regional anaesthesia is not formally contraindicated and general anaesthesia has been used in several surgical procedures. Although there are muscular changes (increased muscle tone and myoclonus) in these patients, the use of non-depolarizing neuromuscular blockers (NMBs) for endotracheal intubation may be very helpful, provided the possibility of abnormal pharmacokinetics and interaction with other drugs is being considered. As most patients suffer from epilepsy, anaesthetists should be aware of the potential risk of seizures associated with volatile anaesthetics. All opioids (alfentanil, fentanyl and remifentanil) activated epileptic discharges and had a dose-related effect on spike activity.

Particular or additional monitoring

Patients with NCL may have a lower baseline body temperature and be at risk for significant hypothermia during general anaesthesia. Adequate monitoring should prevent the complication of profound hypothermia.

Possible complications

The main complications that must be ruled out are aspiration, hypothermia, bradycardia and status epilepticus.

Postoperative care

Seizures are more common in the postoperative period. They may be precipitated by the use of proconvulsant anaesthetic agents, hypoxia, hypercapnoea, electrolyte disturbances (hyponatraemia, hypocalcaemia and hypomagnesaemia), hypoglycaemia, uraemia, subtherapeutic levels of AEDs or local anaesthetic toxicity. Correction of any precipitating factors and checking plasma levels of AEDs are required. It is essential to restart the antiepileptic treatment in the postoperative period as soon as possible.

Additionally, avoiding prolonged mechanical ventilation and temperature control must be among the main goals in the postoperative period. Non-invasive postoperative ventilation may be needed in this period.

Information about emergency-like situations / Differential diagnostics

Caused by the illness to give a tool to distinguish between a side effect of the anaesthetic procedure and a manifestation of the disease.

Not reported.

Ambulatory anaesthesia

Ambulatory anaesthesia, in view of significant risks associated with the severe neurological disease that tend to exacerbate during the postoperative period, can usually not be recommended.

Obstetrical anaesthesia

Not applicable.

Literature and internet links

1. Mole SE, Williams R, Goebel HH, et al. The Neuronal Ceroid Lipofuscinoses (Batten Disease). 2nd ed. Contemporary Neurology Series 2011;Oxford University Press: Oxford 480
2. Schulz A, Kohlschütter A, Mink J, Simonati A, Williams R. NCL diseases - clinical perspectives. *Biochim Biophys Acta* 2013;1832(11):1801-6
3. Kohlschütter A, Schulz A, Denecke J. Epilepsy in Neuronal Ceroid Lipofuscinoses (NCL). *J Pediatr Epilepsy* 2014;3:199-206
4. Miao N, Levin SW, Baker EH, Caruso RC, Zhang Z, Gropman A, Koziol D, Wesley R, Mukherjee AB, Quezado ZM. Children with infantile neuronal ceroid lipofuscinosis have an increased risk of hypothermia and bradycardia during anesthesia. *Anesth Analg* 2009;109(2): 372-8.
5. Pereira D, Pereira M, Caldas F. Anesthesia management in neuronal ceroid lipofuscinoses. *Paediatr Anaesth* 2006;16(3):356-8
6. Yamada Y, Doi K, Sakura S, Saito Y. Anesthetic management for a patient with Jansky-Bielschowsky disease. *Can J Anaesth* 2002;49(1):81-3
7. Johannsen J, Nickel M, Schulz A, Denecke J. Considering Valproate as a Risk Factor for Rapid Exacerbation of Complex Movement Disorder in Progressed Stages of Late-Infantile CLN2 Disease. *Neuropediatrics* 2016
8. <http://www.ncl-net.de/en/index.htm>

Last date of modification: June 2016

These guidelines have been prepared by:

Author

Freda Richa, Anaesthesiologist, Saint-Joseph University, Hotel-Dieu de France Hospital
Beirut, Lebanon
fredrich24@yahoo.com

Peer revision 1

Ruth E. Williams, Consultant Paediatric Neurologist, Department of Paediatric Neurology
The Evelina Children's Hospital, Guy's and St Thomas' NHS Foundation Trust
London, United Kingdom
ruth.williams@gstt.nhs.uk

Peer revision 2

Alfried Kohlschütter, Working Group of degenerative brain diseases, Children's Hospital,
UKE – University Hospital Hamburg-Eppendorf, Germany
kohlschuetter@uke.uni-hamburg.de

Please note that this guideline has not been reviewed by an anaesthesiologist but by two disease experts instead.

Herausgeber



DGAI
Deutsche Gesellschaft
für Anästhesiologie und
Intensivmedizin e.V.
Präsident: Prof. Dr.
B. Zwißler, München



BDA
Berufsverband Deutscher
Anästhesisten e.V.
Präsident: Prof. Dr.
G. Geldner, Ludwigsburg



DAAF
Deutsche Akademie
für Anästhesiologische
Fortbildung e.V.
Präsident: Prof. Dr.
F. Wappler, Köln

Schriftleitung

Präsident/in der Herausgeberverbände
Gesamtschriftleiter:
Prof. Dr. Dr. Kai Zacharowski, Frankfurt
Stellvertretender Gesamtschriftleiter:
Prof. Dr. T. Volk, Homburg/Saar
CME-Schriftleiter:
Prof. Dr. H. A. Adams, Trier

Redaktionskomitee

Prof. Dr. G. Beck, Wiesbaden
Dr. iur. E. Biermann, Nürnberg
Prof. Dr. H. Bürkle, Freiburg
Prof. Dr. B. Ellger, Münster
Prof. Dr. K. Engelhard, Mainz
Prof. Dr. M. Fischer, Göppingen
Priv.-Doz. Dr. T. Iber, Baden-Baden
Prof. Dr. U. X. Kaisers, Ulm
Prof. Dr. W. Meißner, Jena
Prof. Dr. C. Nau, Lübeck
Dr. M. Rähler, Mainz
Prof. Dr. A. Schleppers, Nürnberg
Prof. Dr. G. Theilmeier, Hannover
Prof. Dr. M. Thiel, Mannheim
Prof. Dr. F. Wappler, Köln
Prof. Dr. M. Weigand, Heidelberg

Redaktion

Alexandra Hisom M.A. &
Dipl.-Sozw. Holger Sorgatz
Korrespondenzadresse: Roritzerstraße 27 |
90419 Nürnberg | Deutschland
Tel.: 0911 9337812 | Fax: 0911 3938195
E-Mail: anaesth.intensivmed@dgai-ev.de

Verlag & Druckerei

Aktiv Druck & Verlag GmbH
An der Lohwiese 36 |
97500 Ebelsbach | Deutschland
www.aktiv-druck.de

Geschäftsführung

Wolfgang Schröder | Jan Schröder |
Nadja Schwarz
Tel.: 09522 943560 | Fax: 09522 943567
E-Mail: info@aktiv-druck.de

Anzeigen | Vertrieb

Pia Engelhardt
Tel.: 09522 943570 | Fax: 09522 943577
E-Mail: anzeigen@aktiv-druck.de

Verlagsrepräsentanz

Rosi Braun
PF 13 02 26 | 64242 Darmstadt
Tel.: 06151 54660 | Fax: 06151 595617
E-Mail: rbraunwerb@aol.com

Herstellung | Gestaltung

Manfred Wuttke | Stefanie Triebert
Tel.: 09522 943571 | Fax: 09522 943577
E-Mail: ai@aktiv-druck.de

Titelbild

Dipl.-Designerin Monique Minde,
Nürnberg

Erscheinungsweise 2017

Der 58. Jahrgang erscheint jeweils zum
Monatsanfang, Heft 7/8 als Doppelausgabe.

Bezugspreise (inkl. Versandkosten):

- **Einzelhefte** 30,- €
- **Jahresabonnement:**
 - Europa (ohne Schweiz) 258,- €
(inkl. 7 % MwSt.)
 - Schweiz 266,- €
 - Rest der Welt 241,- €

Mitarbeiter aus Pflege, Labor, Studenten und Auszubildende (bei Vorlage eines entsprechenden Nachweises)

- Europa (ohne Schweiz) 94,- €
(inkl. 7 % MwSt.)
- Schweiz 90,- €
- Rest der Welt 94,- €

**Für Mitglieder der DGAI und/oder
des BDA ist der Bezug der Zeitschrift
im Mitgliedsbeitrag enthalten.**

Allgemeine Geschäfts- und Liefer- bedingungen

Die allgemeinen Geschäfts- und Liefer-
bedingungen entnehmen Sie bitte dem
Impressum auf www.ai-online.info

Indexed in **Current Contents®/Clinical
Medicine, EMBASE/Excerpta Medica;
Medical Documentation Service;
Research Alert; Sci Search; SUBIS
Current Awareness in Biomedicine;
VINITI: Russian Academy of Science.**

Nachdruck | Urheberrecht

Die veröffentlichten Beiträge sind urhe-
berrechtlich geschützt. Jegliche Art von
Vervielfältigungen – sei es auf mechani-
schem, digitalem oder sonst möglichem
Wege – bleibt vorbehalten. Die Aktiv
Druck & Verlags GmbH ist allein auto-
risiert, Rechte zu vergeben und Sonder-
drucke für gewerbliche Zwecke, gleich
in welcher Sprache, herzustellen. An-
fragen hierzu sind nur an den Verlag zu
richten. Jede im Bereich eines gewerbli-
chen Unternehmens zulässig hergestellte
oder benutzte Kopie dient gewerblichen
Zwecken gem. § 54 (2) UrhG. Die Wie-
dergabe von Gebrauchsnamen, Handels-
namen, Warenbezeichnungen usw. in
dieser Zeitschrift berechtigt auch ohne
besondere Kennzeichnung nicht zu der
Annahme, dass solche Namen im Sinne
der Warenzeichen- und Markenschutz-
Gesetzgebung als frei zu betrachten wä-
ren und daher von jedermann benutzt
werden dürften.

Wichtiger Hinweis

Für Angaben über Dosierungsanwei-
sungen und Applikationsformen kann
vom Verlag und den Herausgebern keine
Gewähr übernommen werden. Derartige
Angaben müssen vom jeweiligen An-
wender im Einzelfall anhand anderer
Literaturstellen auf ihre Richtigkeit über-
prüft werden. Gleiches gilt für berufs-
und verbandspolitische Stellungnahmen
und Empfehlungen.

Online-Ausgabe der A&I ab April 2017 open access: www.ai-online.info

CONTACT US

Please do not hesitate to contact us. We will be glad to answer and provide further information to you at any time.

.....
Name

.....
First Name

.....
Department / Hospital

.....
Place

.....
Telephone

.....
E-Mail

.....
Date / Signature

Please contact me for further information

I would like to participate in the project

ADDRESS

German Society of Anaesthesiology and
Intensive Care Medicine
Nina Schnabel
Roritzerstrasse 27 | 90419 Nuremberg | Germany
Tel.: +49-911-9337822 | Fax: +49-911-3938195
Email: nschnabel@orphananesthesia.eu