

A&I

ANÄSTHESIOLOGIE & INTENSIVMEDIZIN

Offizielles Organ: Deutsche Gesellschaft für Anästhesiologie und Intensivmedizin e.V. (DGAI)
Berufsverband Deutscher Anästhesisten e.V. (BDA)
Deutsche Akademie für Anästhesiologische Fortbildung e.V. (DAAF)
Organ: Deutsche Interdisziplinäre Vereinigung für Intensiv- und Notfallmedizin e.V. (DIVI)



Rett syndrome

ROHHAD syndrome

orphan**a**nesthesia

a project of the German Society
of Anaesthesiology and Intensive Care Medicine

SUPPLEMENT NR. 6 | 2018

OrphanAnesthesia –

ein krankheitsübergreifendes Projekt des Wissenschaftlichen Arbeitskreises Kinderanästhesie der Deutschen Gesellschaft für Anästhesiologie und Intensivmedizin e.V.

Ziel des Projektes ist die Veröffentlichung von Handlungsempfehlungen zur anästhesiologischen Betreuung von Patienten mit seltenen Erkrankungen. Damit will OrphanAnesthesia einen wichtigen Beitrag zur Erhöhung der Patientensicherheit leisten.

Patienten mit seltenen Erkrankungen benötigen für verschiedene diagnostische oder therapeutische Prozeduren eine anästhesiologische Betreuung, die mit einem erhöhten Risiko für anästhesieassoziierte Komplikationen einhergehen. Weil diese Erkrankungen selten auftreten, können Anästhesisten damit keine Erfahrungen gesammelt haben, so dass für die Planung der Narkose die Einholung weiterer Information unerlässlich ist. Durch vorhandene spezifische Informationen kann die Inzidenz von mit der Narkose assoziierten Komplikationen gesenkt werden. Zur Verfügung stehendes Wissen schafft Sicherheit im Prozess der Patientenversorgung.

Die Handlungsempfehlungen von OrphanAnesthesia sind standardisiert und durchlaufen nach ihrer Erstellung einen Peer-Review-Prozess, an dem ein Anästhesist sowie ein weiterer Krankheitsexperte (z.B. Pädiater oder Neurologe) beteiligt sind. Das Projekt ist international ausgerichtet, so dass die Handlungsempfehlungen grundsätzlich in englischer Sprache veröffentlicht werden.

Ab Heft 5/2014 werden im monatlichen Rhythmus je zwei Handlungsempfehlungen als Supplement der A&I unter www.ai-online.info veröffentlicht. Als Bestandteil der A&I sind die Handlungsempfehlungen damit auch zitierfähig. Sonderdrucke können gegen Entgelt bestellt werden.

OrphanAnesthesia –

a common project of the Scientific Working Group of Paediatric Anaesthesia of the German Society of Anaesthesiology and Intensive Care Medicine

The target of OrphanAnesthesia is the publication of anaesthesia recommendations for patients suffering from rare diseases in order to improve patients' safety. When it comes to the management of patients with rare diseases, there are only sparse evidence-based facts and even far less knowledge in the anaesthetic outcome. OrphanAnesthesia would like to merge this knowledge based on scientific publications and proven experience of specialists making it available for physicians worldwide free of charge.

All OrphanAnesthesia recommendations are standardized and need to pass a peer review process. They are being reviewed by at least one anaesthesiologist and another disease expert (e.g. paediatrician or neurologist) involved in the treatment of this group of patients.

The project OrphanAnesthesia is internationally oriented. Thus all recommendations will be published in English.

Starting with issue 5/2014, we'll publish the OrphanAnesthesia recommendations as a monthly supplement of A&I (Anästhesiologie & Intensivmedizin). Thus they can be accessed and downloaded via www.ai-online.info. As being part of the journal, the recommendations will be quotable. Reprints can be ordered for payment.

**Bisher in A&I publizierte
Handlungsempfehlungen finden
Sie unter:**

www.ai-online.info/Orphsuppl
www.orphananesthesia.eu

**A survey of until now in A&I
published guidelines can be
found on:**

www.ai-online.info/Orphsuppl
www.orphananesthesia.eu

Projektleitung

Prof. Dr. Tino Münster, MHBA
Geschäftsführender Oberarzt
Facharzt für Anästhesie,
Spezielle Schmerztherapie,
Notfallmedizin
Anästhesiologische Klinik
Friedrich-Alexander-Universität
Erlangen-Nürnberg
Krankenhausstraße 12
91054 Erlangen, Deutschland
Tel.: 09131 8542441
Fax: 09131 8536147
E-Mail: muenster@kfa.imed.uni-erlangen.de

orphananesthesia

Anaesthesia recommendations for patients suffering from Rett syndrome

Disease name: Rett syndrome (RS)

ICD 10: F84.2

Synonyms: Autistic disorder (F84.0)

Rett syndrome is a rare inherited neurodevelopmental disease originally described by Andreas Rett in 1966, which occurs mostly in girls [1] and causes developmental and nervous system problems. Genetically, Rett syndrome results from mutations in the genes encoding methyl-cytosine-guanosine (CpG) binding protein 2 (MECP2) located on the X chromosome. In less than 10% of RTT cases, mutations in the genes CDKL5 or FOXP1 have also been found to resemble it [2]. Affected individuals have clinical features that overlap with autism. Infants with Rett syndrome seem to grow and develop normally in the early months. However, between 6 and 18 months and 3 years of age development stagnates and they begin to lose previously attained skills.

Medicine in progress



Perhaps new knowledge

Every patient is unique

Perhaps the diagnosis is wrong

► **Citation:** Maheshwari N: Rett syndrome. Anästh Intensivmed 2018;59:S117-S124.
DOI: 10.19224/ai2018.S117

Disease summary

Symptoms include

- loss of speech (partial or complete)
- loss of purposeful hand movements such as grasping (partial or complete)
- stereotypic hand movements such as hand wringing or hand-washing
- balance problems (ataxia/apraxia of gait)
- breathing problems (periodic breathing with hyperventilation and breath holding)
- behaviour problems (particularly anxiety, self-abusive or aggressive)
- intellectual disability

The clinical criteria for the diagnosis of classic Rett syndrome were developed initially in the 1980s and recently updated in 2010. These criteria include an apparently normal pre- or perinatal period and first 6 months of life. Over the following months, there is a loss of developmental skills, which progresses to severe mental retardation and characteristic hand-wringing movements [3,4]. Other features include autistic-like behaviours, oral-motor dysfunctions, gastrointestinal motility disorders (including abdominal bloating), scoliosis, autonomic dysfunction and somatic developmental impairment [5]. Various movement disorders are exhibited including dystonia, choreoathetosis and myoclonic jerks. [6,7]

Rett syndrome has no cure. We can treat some of the symptoms with medication, surgery, and physical, occupational and speech therapy. Most people with RS live into middle age and beyond. They will usually need care throughout their lives.

Typical surgery

Scoliosis is common in RS and increases in frequency with age through puberty [8]. Scoliosis in individuals with Rett syndrome often does not respond to orthotics. Surgery should be considered in patients with Cobb angles of more than 40-50° or curves that cause pain or loss of function, and is best performed with a multidisciplinary team providing specialist anaesthetic and pain management support, seizure control, with the aim of promoting early mobilisation [9]. Surgeons usually try to wait until the child is at least 10 years of age before performing spinal fusion. They may often recommend bracing, physiotherapy and physical activity to delay surgery in order to allow for as much growth as possible. The aims of surgery are to achieve a balanced, fused spine so that the shoulders and hips are more level in upright positions, such as sitting or standing.

Surgery may be indicated for other reasons than orthopaedic issues, e.g. tonsillectomy and adenoidectomy for the treatment of obstructive sleep apnoea (OSA), placement of a gastrostomy in case of feeding difficulties, hip subluxation or possibly tendon release due to significant contractures.

Dental procedures.

Type of anaesthesia

An altered muscle tone and thoracic deformity might complicate the administration of general anaesthesia [10]. Patients with Rett syndrome are at increased risk of sedation-related transient adverse events. Most adverse events are related to prolonged apnoea and airway compromise. Patients with Rett syndrome should not be sedated without appropriate personnel available who are skilled to intervene and recognise the risks involved in sedating this unique population [11]. In patients with a genuine aspiration risk, rapid sequence intubation (RSI) with cricoid pressure may be considered.

As these patients are unable to cooperate and have continuous abnormal limb movements, spinal or epidural anaesthesia is difficult. In addition, scoliosis could make placement of an epidural catheter difficult.

Necessary additional diagnostic procedures (preoperative)

Preoperative evaluation of a patient with RS is very important. We should check:

- the breathing patterns,
- arterial blood gases and lactic acid levels preoperatively (although arterial blood gases may be affected, lactic acid levels are rarely increased),
- the presence of gastrointestinal reflux and autonomic disturbances,
- EEG for seizure activity (could be a concern in pre or postoperative periods),
- ECG for arrhythmias associated with prolonged QT syndrome, and
- the musculoskeletal system.

Parents and caregivers should be consulted about any other relevant information about the patient.

Particular preparation for airway management

A primary concern for anaesthesia providers is the potential for difficult endotracheal intubation related to limited mouth opening, micrognathia, and limited neck movement. As such, the ability to accomplish adequate bag-valve mask ventilation should be demonstrated prior to the use of neuromuscular blocking agents. Additionally, appropriate equipment for dealing with the difficult airway should be readily available, including indirect laryngoscopy tools or a tracheostomy set to secure the airway in the event of severe difficulty when securing the airway.

Various options for airway management in patients with limited mouth opening like antegrade fibre-optic-guided orotracheal or nasotracheal intubation, retrograde guidewire-assisted fibre-optic intubation and blind nasal intubation may be considered.

Particular preparation for transfusion or administration of blood products

Not reported.

Particular preparation for anticoagulation

Not reported.

Particular precautions for positioning, transport or mobilisation

Not reported.

Probable interaction between anaesthetic agents and patient's long-term medication

Not reported.

Anaesthesiologic procedure

- Inhalational and intravenous anaesthetic agents are generally considered to have potent anticonvulsant properties. Many of these agents, including the barbiturates, propofol, and the volatile agents, have been used successfully.
- As the residual effects of anaesthetic agents may impact on both upper airway control and postoperative respiratory function, whenever feasible, short-acting agents whose effects dissipate rapidly, such as remifentanyl, should be considered.
- Use of thiopental and succinylcholine, prokinetics such as cisapride, antipsychotics such as thioridazine, tricyclic antidepressants such as imipramine, antiarrhythmics such as quinidine amiodarone, sotalol as well as erythromycin and ketoconazole should be avoided because of the potential to provoke QT abnormalities and cardiac arrhythmias.
- These patients are excessively sensitive to sedative drugs, and delayed recovery from anaesthesia has been reported. Hence, the drugs should be titrated to maintain an optimal depth of anaesthesia and avoid overdose. Spontaneous ventilation rather than controlled ventilation is preferred, as this would require administration of less anaesthetic agents.
- Additionally, anatomical malposition of the vessels may occur in these patients. The use of ultrasound guidance may be invaluable to aid in gaining adequate vascular access for major surgical procedures [12].

Particular or additional monitoring

Anaesthetic considerations for patients with a prolonged QTc interval include avoidance of potential factors that may further prolong the QT interval, including an increase in sympathetic tone; hypothermia; electrolyte abnormalities including hypokalaemia, hypomagnesemia, and hypocalcaemia; as well as various medications which may be administered during the perioperative period, as mentioned above.

Possible complications

- Hypermetabolic states observed in patients with Rett syndrome raise the possibility that anaesthesia and anaesthetic agents may increase the risk of malignant hyperthermia, although this has not been reported in patients.
- The use of succinylcholine is controversial because of possible hyperkalaemia and prolongation of the QTc interval, which may predispose the patient to fatal arrhythmias.

Postoperative care

- Regardless of the agents used, given the airway and respiratory co-morbid conditions seen in Rett syndrome, postoperative monitoring of respiratory function is recommended. Although postoperative mechanical ventilation can be anticipated in these patients, consideration must be given to the antagonism of neuromuscular blockade and the reversal of anaesthesia when circumstances so dictate.
- Previous anecdotal reports of anaesthesia in children with Rett syndrome have noted postoperative respiratory problems, including atelectasis and respiratory infections, which may require postoperative mechanical ventilation or reintubation.
- Non-invasive respiratory support techniques such as BiPAP would facilitate postoperative tracheal extubation in these patients.

Information about emergency-like situations / differential diagnostics

... [...] ... caused by the illness to give a tool to distinguish between a side effect of the anaesthetic procedure and a manifestation of the disease

The risk of prolonged QT interval and T-wave changes increases with age. These ECG abnormalities are considered as potential causes of sudden death encountered in up to one-third of patients with Rett syndrome. Aside from medications, other intraoperative events may lead to a further prolongation of the QT interval, including increased sympathetic tone, which has been demonstrated to occur during endotracheal intubation. To minimise the risk of this, an adequate depth of anaesthesia should be ensured during such procedures.

Ambulatory anaesthesia

Ambulatory anaesthesia is possible following the commonly accepted guidelines if the procedure itself does not require a longer phase of supervision. This applies especially to oral surgery.

Obstetrical anaesthesia

Not reported.

Literature and internet links

1. Rett A. On a unusual brain atrophy syndrome in hyperammonemia in childhood. *Wien Med Wochenschr* 1966;116:723-726
2. Amir RE, Van den Veyver IB, Wan M, Tran CQ, Francke U, Zoghbi H. Rett syndrome is caused by mutations in X-linked *MECP2*, encoding methyl-CpG-binding protein 2. *Nature Genet* 1999; 23:185-188
3. Hagberg B, Hanefeld F, Percy AK, Skjeldal. An update on clinically applicable diagnostic criteria in Rett syndrome. *Eur J Paediatr Neurol* 2002;6:293-297
4. Neul JL, Kaufmann WE, Glaze DG, Christodoulou J, Clarke AJ, Bahi-Buisson N, Leonard H, Bailey MES, Schanen NC, Zappella M, Renieri A, Huppke P, Percy A, for the RettSearch Consortium. Rett Syndrome: Revised Diagnostic Criteria and Nomenclature. *Ann Neurol* 2010;68:946-951
5. Armstrong DD. Review of Rett syndrome. *J Neuropathol Exp Neurol* 1997;56(8):843-849
6. Al-Mateen M, Philippart M, Shields WD. Rett syndrome: A commonly overlooked progressive encephalopathy in girls. *Am J Dis Child* 1986;140(8):761-765
7. Hagberg B, Aicardi J, Dias K, Ramos O. A progressive syndrome of autism, dementia, ataxia and loss of purposeful hand use in girls: Rett's syndrome: report of 35 cases. *Ann Neurol* 1983;14(4):471-479
8. Percy AK, Lane JB, Childers J, Skinner S, Annese F, Barrish J, Caeg E, Glaze DG, MacLeod P. Rett syndrome: North American database. *J Child Neurol* 2007;22:1338-1341
9. Downs J, Bergman A, Carter P, Anderson A, Palmer GM, Roye D, van Bosse H, Bebbington A, Larsson EL, Smith BG, Baikie G, Fyfe S, Leonard H. Guidelines for management of scoliosis in Rett syndrome patients based on expert consensus and clinical evidence. *Spine (Phila Pa 1976)*. 2009;1;34(17)
10. Lugaresi E, Girignotta F, Montagna P. Breathing in Rett syndrome. *Brain Dev* 1985;7:329-333
11. Tofil N, Buckmaster M, Callans B, Winkler M, Islam M, Percy A. Deep sedation with Propofol in patients with Rett syndrome. *J Child Neurol* 2006;21:210-213
12. Doniger SJ, Ishimine P, Fox JC, Kanegaye JT. Randomized controlled trial of ultrasoundguided peripheral intravenous catheter placement versus traditional techniques in difficult access pediatric patients. *Pediatr Emerg Care* 2009;25:154-159.

Last date of modification: August 2015

These guidelines have been prepared by:

Author

Namrata Maheshwari, Anaesthesiologist, Department of Critical Care, Fortis Hospital, Mohali, India
dnamratamaheshwari@gmail.com

Peer revision 1

John Christodoulou, Director, Western Sydney Genetics Program, Head Discipline of Genetic Medicine, Sydney Medical School, University of Sydney, Australia
johnc@chw.edu.au

Peer revision 2

Alan Percy, Department of Neurobiology, Civitan International Research Center, University of Alabama at Birmingham, USA
apercy@uab.edu

Please note that this guideline has not been reviewed by an anaesthesiologist but by two disease experts instead.

Herausgeber



DGAI

Deutsche Gesellschaft
für Anästhesiologie und
Intensivmedizin e.V.
Präsident: Prof. Dr.
B. Zwißler, München



BDA

Berufsverband Deutscher
Anästhesisten e.V.
Präsident: Prof. Dr.
G. Geldner, Ludwigsburg



DAAF

Deutsche Akademie
für Anästhesiologische
Fortbildung e.V.
Präsident: Prof. Dr.
F. Wappler, Köln

Schriftleitung

Präsident/in der Herausgeberverbände
Gesamtschriftleiter:
Prof. Dr. Dr. Kai Zacharowski, Frankfurt
Stellvertretender Gesamtschriftleiter:
Prof. Dr. T. Volk, Homburg/Saar
CME-Schriftleiter:
Prof. Dr. H. A. Adams, Trier

Redaktionskomitee

Prof. Dr. G. Beck, Wiesbaden
Dr. iur. E. Biermann, Nürnberg
Prof. Dr. H. Bürkle, Freiburg
Prof. Dr. B. Ellger, Dortmund
Prof. Dr. K. Engelhard, Mainz
Prof. Dr. M. Fischer, Göppingen
Priv.-Doz. Dr. T. Iber, Baden-Baden
Prof. Dr. U. X. Kaisers, Ulm
Prof. Dr. W. Meißner, Jena
Prof. Dr. C. Nau, Lübeck
Dr. M. Rähmer, Mainz
Prof. Dr. A. Schleppers, Nürnberg
Prof. Dr. G. Theilmeier, Hannover
Prof. Dr. M. Thiel, Mannheim
Prof. Dr. F. Wappler, Köln
Prof. Dr. M. Weigand, Heidelberg

Redaktion

Alexandra Hisom M.A. &
Dipl.-Sozw. Holger Sorgatz
Korrespondenzadresse: Roritzerstraße 27 |
90419 Nürnberg | Deutschland
Tel.: 0911 9337812 | Fax: 0911 3938195
E-Mail: anaesth.intensivmed@dgai-ev.de

Verlag & Druckerei

Aktiv Druck & Verlag GmbH

An der Lohwiese 36 |
97500 Ebelsbach | Deutschland
www.aktiv-druck.de

Geschäftsführung

Wolfgang Schröder | Jan Schröder |
Nadja Schwarz
Tel.: 09522 943560 | Fax: 09522 943567
E-Mail: info@aktiv-druck.de

Anzeigen | Vertrieb

Pia Engelhardt
Tel.: 09522 943570 | Fax: 09522 943577
E-Mail: anzeigen@aktiv-druck.de

Verlagsrepräsentanz

Jürgen Distler
Roritzerstraße 27, 90419 Nürnberg
Tel.: 0171 9432534 | Fax: 0911 3938195
E-Mail: jdistler@bda-ev.de

Herstellung | Gestaltung

Manfred Wuttke | Stefanie Triebert
Tel.: 09522 943571 | Fax: 09522 943577
E-Mail: ai@aktiv-druck.de

Titelbild

Dipl.-Designerin Monique Minde,
Nürnberg

Erscheinungsweise 2018

Der 59. Jahrgang erscheint jeweils zum
Monatsanfang, Heft 7/8 als Doppelausgabe.

Bezugspreise (inkl. Versandkosten):

• Einzelhefte	30,- €
• Jahresabonnement:	
Europa (ohne Schweiz)	258,- €
(inkl. 7 % MwSt.)	
Schweiz	266,- €
Rest der Welt	241,- €

Mitarbeiter aus Pflege, Labor, Studenten und Auszubildende (bei Vorlage eines entsprechenden Nachweises)

Europa (ohne Schweiz)	94,- €
(inkl. 7 % MwSt.)	
Schweiz	90,- €
Rest der Welt	94,- €

**Für Mitglieder der DGAI und/oder
des BDA ist der Bezug der Zeitschrift
im Mitgliedsbeitrag enthalten.**

Allgemeine Geschäfts- und Liefer- bedingungen

Die allgemeinen Geschäfts- und Liefer-
bedingungen entnehmen Sie bitte dem
Impressum auf www.ai-online.info

Indexed in **Current Contents®/Clinical
Medicine, EMBASE/Excerpta Medica;
Medical Documentation Service;
Research Alert; Sci Search; SUBIS
Current Awareness in Biomedicine;
VINITI: Russian Academy of Science.**

Nachdruck | Urheberrecht

Die veröffentlichten Beiträge sind urhe-
berrechtlich geschützt. Jegliche Art von
Vervielfältigungen – sei es auf mechani-
schem, digitalem oder sonst möglichem
Wege – bleibt vorbehalten. Die Aktiv
Druck & Verlags GmbH ist allein auto-
risiert, Rechte zu vergeben und Sonder-
drucke für gewerbliche Zwecke, gleich
in welcher Sprache, herzustellen. An-
fragen hierzu sind nur an den Verlag zu
richten. Jede im Bereich eines gewerbli-
chen Unternehmens zulässig hergestellte
oder benutzte Kopie dient gewerblichen
Zwecken gem. § 54 (2) UrhG. Die Wie-
dergabe von Gebrauchsnamen, Handels-
namen, Warenbezeichnungen usw. in
dieser Zeitschrift berechtigt auch ohne
besondere Kennzeichnung nicht zu der
Annahme, dass solche Namen im Sinne
der Warenzeichen- und Markenschutz-
Gesetzgebung als frei zu betrachten wä-
ren und daher von jedermann benutzt
werden dürften.

Wichtiger Hinweis

Für Angaben über Dosierungsanwei-
sungen und Applikationsformen kann
vom Verlag und den Herausgebern keine
Gewähr übernommen werden. Derartige
Angaben müssen vom jeweiligen An-
wender im Einzelfall anhand anderer
Literaturstellen auf ihre Richtigkeit über-
prüft werden. Gleiches gilt für berufs-
und verbandspolitische Stellungnahmen
und Empfehlungen.

Online-Ausgabe der A&I ab April 2017 open access: www.ai-online.info

CONTACT US

Please do not hesitate to contact us. We will be glad to answer and provide further information to you at any time.

.....
Name

.....
First Name

.....
Department / Hospital

.....
Place

.....
Telephone

.....
E-Mail

.....
Date / Signature

Please contact me for further information

☐

I would like to participate in the project

☐

ADDRESS

German Society of Anaesthesiology and
Intensive Care Medicine
Nina Schnabel
Roritzerstrasse 27 | 90419 Nuremberg | Germany
Tel.: +49-911-9337822 | Fax: +49-911-3938195
Email: nschnabel@orphananesthesia.eu