

A&I

ANÄSTHESIOLOGIE & INTENSIVMEDIZIN

Offizielles Organ: Deutsche Gesellschaft für Anästhesiologie und Intensivmedizin e.V. (DGAI)
Berufsverband Deutscher Anästhesisten e.V. (BDA)
Deutsche Akademie für Anästhesiologische Fortbildung e.V. (DAAF)
Organ: Deutsche Interdisziplinäre Vereinigung für Intensiv- und Notfallmedizin e.V. (DIVI)



Trisomy 18

VACTERL association

orphan**a**nesthesia

a project of the German Society
of Anaesthesiology and Intensive Care Medicine

SUPPLEMENT NR. 4 | 2019

OrphanAnesthesia –

ein krankheitsübergreifendes Projekt des Wissenschaftlichen Arbeitskreises Kinderanästhesie der Deutschen Gesellschaft für Anästhesiologie und Intensivmedizin e.V.

Ziel des Projektes ist die Veröffentlichung von Handlungsempfehlungen zur anästhesiologischen Betreuung von Patienten mit seltenen Erkrankungen. Damit will Orphan Anesthesia einen wichtigen Beitrag zur Erhöhung der Patientensicherheit leisten.

Patienten mit seltenen Erkrankungen benötigen für verschiedene diagnostische oder therapeutische Prozeduren eine anästhesiologische Betreuung, die mit einem erhöhten Risiko für anästhesieassoziierte Komplikationen einhergehen. Weil diese Erkrankungen selten auftreten, können Anästhesisten damit keine Erfahrungen gesammelt haben, so dass für die Planung der Narkose die Einholung weiterer Information unerlässlich ist. Durch vorhandene spezifische Informationen kann die Inzidenz von mit der Narkose assoziierten Komplikationen gesenkt werden. Zur Verfügung stehendes Wissen schafft Sicherheit im Prozess der Patientenversorgung.

Die Handlungsempfehlungen von OrphanAnesthesia sind standardisiert und durchlaufen nach ihrer Erstellung einen Peer-Review-Prozess, an dem ein Anästhesist sowie ein weiterer Krankheitsexperte (z.B. Pädiater oder Neurologe) beteiligt sind. Das Projekt ist international ausgerichtet, so dass die Handlungsempfehlungen grundsätzlich in englischer Sprache veröffentlicht werden.

Ab Heft 5/2014 werden im monatlichen Rhythmus je zwei Handlungsempfehlungen als Supplement der A&I unter www.ai-online.info veröffentlicht. Als Bestandteil der A&I sind die Handlungsempfehlungen damit auch zitierfähig. Sonderdrucke können gegen Entgelt bestellt werden.

OrphanAnesthesia –

a common project of the Scientific Working Group of Paediatric Anaesthesia of the German Society of Anaesthesiology and Intensive Care Medicine

The target of OrphanAnesthesia is the publication of anaesthesia recommendations for patients suffering from rare diseases in order to improve patients' safety. When it comes to the management of patients with rare diseases, there are only sparse evidence-based facts and even far less knowledge in the anaesthetic outcome. OrphanAnesthesia would like to merge this knowledge based on scientific publications and proven experience of specialists making it available for physicians worldwide free of charge.

All OrphanAnesthesia recommendations are standardized and need to pass a peer review process. They are being reviewed by at least one anaesthesiologist and another disease expert (e.g. paediatrician or neurologist) involved in the treatment of this group of patients.

The project OrphanAnesthesia is internationally oriented. Thus all recommendations will be published in English.

Starting with issue 5/2014, we'll publish the OrphanAnesthesia recommendations as a monthly supplement of A&I (Anästhesiologie & Intensivmedizin). Thus they can be accessed and downloaded via www.ai-online.info. As being part of the journal, the recommendations will be quotable. Reprints can be ordered for payment.

**Bisher in A&I publizierte
Handlungsempfehlungen finden
Sie unter:**

www.ai-online.info/Orphsuppl
www.orphananesthesia.eu

**A survey of until now in A&I
published guidelines can be
found on:**

www.ai-online.info/Orphsuppl
www.orphananesthesia.eu

Projektleitung

Prof. Dr. Tino Münster, MHBA

Chefarzt
Klinik für Anästhesie und
operative Intensivmedizin
Krankenhaus Barmherzige
Brüder Regensburg
Prüfener Straße 86
93049 Regensburg,
Deutschland

Tel.: 0941 369-2350

E-Mail: Tino.Muenster@barmherzige-regensburg.de

orphananesthesia

Anaesthesia recommendations for VACTERL association

Disease name: VACTERL association

ICD 10: Q87.2

Synonyms: VATERS association, VACTERLS association, VACTERL association (ORPHA887), VATER association (ORPHA887), VATER syndrome, ORPHA887. Each letter of the mnemonic stands for one or more type of malformation. It is an association rather than a syndrome, as there is no evidence that the malformations are pathogenetically related. However, they occur together more frequently than expected by chance.

VACTERL association is defined by the presence of at least 3 of the following congenital malformations: vertebral defects, anal atresia, cardiac defects, tracheoesophageal fistula, renal anomalies and limb abnormalities. In addition to these core component features, patients may also have other congenital anomalies.

Medicine in progress



Perhaps new knowledge

Every patient is unique

Perhaps the diagnosis is wrong



Find more information on the disease, its centres of reference and patient organisations on Orphanet: www.orpha.net

► **Citation:** Richards E: VACTERL association. Anästh Intensivmed 2019;60:S83–S94.
DOI: 10.19224/ai2019.S083

Disease summary

Exact incidence unknown but in the order of 1/10,000 -1/40,000 live born infants.

Originally described in 1973 by Quan and Smith, as VATER association (acronym standing for **V**erterbral anomalies, **A**nal atresia, **T**racheoesophageal fistula with **O**esophageal atresia, **R**adial and **R**enal dysplasia). In 1974, Temtamy and Miller included ventricular septal defect and single umbilical artery to the 'V'. But in 1975, this was changed to VACTERL by Kaufman and Nora and Nora (where 'C' stood for **C**ardiac anomalies and 'L' included limb rather than just radial anomalies).

VACTERL association is a sporadic disease and a positive family history requires careful differential diagnosis with other genetic conditions. Though the exact cause is unknown, it is thought to be multifactorial in aetiology, with environmental triggers, including teratogens, interacting with a genetically susceptible genome. Triggers include exposure of the foetus to sex hormones, anticholesterol drugs, lead, adriamycin and dibenzepin in the first trimester, as well as babies born to diabetic mothers. It is rarely seen more than once in one family.

Due to the number of organs affected non-randomly, it is thought that a "developmental field defect" occurs during blastogenesis (2–4 weeks of gestation), where abnormal structures are derived from the embryonic mesoderm. Males seem more affected than females and rarely multiple individuals are affected in one family.

VACTERL association is a diagnosis of exclusion. The diagnosis is made clinically when 3 or more congenital defects are present and no clinical or laboratory based evidence exists for the presence of one of the many similar overlapping conditions. No validated diagnostic criteria have been published.

Antenatal diagnosis is challenging, as both skill and experience is needed to interpret scans performed and some features of the association are difficult to detect prior to birth.

Overall prognosis depends on type and severity of anomalies present. Today, the decreased mortality in most children is the result of early detection by ultrasound in the second trimester as well as early surgical intervention and rehabilitation.

Genetic counselling is difficult because of lack of information. Children with VACTERL association have normal development and normal intelligence.

V – Vertebral and Vascular anomalies 70% (60–80%)

Small hypoplastic/dysplastic/missing/supernumerary vertebrae; hemivertebrae; "butterfly" vertebrae; wedge vertebrae; vertebral clefts and fusion; caudal regression; tethered cord; branchial arch/cleft abnormalities; rib anomalies; sacral agenesis; dysplastic sacral vertebrae; scoliosis or kyphoscoliosis secondary to costovertebral anomalies; C5-6 dislocation and severe stenosis with spinal cord impingement, which is rare, and developmental delay with signs of myelopathy which is even rarer.

Early complications – minimal; late complications – risk of developing scoliosis/back pain

Single umbilical artery 20% (often included as part of the 'V' in VACTERL). Antenatal diagnosis of the single umbilical artery is maybe the first sign of the diagnosis but it is not specific of VACTERL association.

A – Anal atresia / imperforate anus 55% (up to 90%)

Noted at birth, requiring surgery in the first days of life, sometimes several surgeries required to fully reconstruct the intestine and anal canal.

Involvement of rectum/anus relates to major risk of genital abnormalities, especially in females with the risk of rectovaginal fistulas and urogenital complications.

C – Cardiovascular anomalies 75% (40–80%)

Most common defects – Ventricular Septal Defect (VSD) (22–30%) +/- heart failure / LV dilatation, Atrial Septal Defect (ASD), Tetralogy of Fallot (TOF)

Less common defects – truncus arteriosus, transposition of the great arteries, hypoplastic left heart syndrome (sporadic reports), patent ductus arteriosus (PDA), coarctation of the aorta

T – Tracheoesophageal fistula / E- Esophageal atresia 32%

15–33% will have associated uncomplicated congenital heart disease, e.g. VSD that do not require surgery.

Oesophageal atresia can occur as an isolated defect with an incidence of around 8%.

R – Renal anomalies 50–80%

Can be severe with incomplete formation of one or both kidneys or urological problems, e.g. severe reflux or obstruction of outflow of urine from the kidneys; Horseshoe kidneys, cystic, aplastic, dysplastic or ectopic kidneys, hydronephrosis; unilateral +/- bilateral agenesis; pyelonephritis; nephrolithiasis.

Kidney failure can occur early in life and may require kidney transplant.

Other renal abnormalities that can occur but are typically considered non-VACTERL include hypospadias, UTI; urethral atresia/stricture, ureteral malformation; genital abnormalities, fistula connecting genitor-urinary (GU) and anorectal tracts (up to 25%).

L – Limb defects incl. radial anomalies up to 70% (40–50%),

Includes displaced, absent or hypoplastic thumb/s, polydactyly, syndactyly and radial aplasia/dysplasia/hypoplasia; radial ray deformities; radioulnar synostosis; club foot; hypoplasia of great toe/tibia; lower limb tibial deformities.

Bilateral limb defects tend to have kidney/urological problems on both sides. Unilateral limb defects tend to have kidney/urological problems on the same side.

Other – Growth deficiencies; failure to thrive.

Typical surgery

VACTERL association defects are treated post birth with issues being approached one at a time.

Management is divided into 2 stages: 1) centers around surgical correction of the specific congenital anomalies, incompatible with life, e.g. tracheoesophageal fistula, certain types of severe cardiac malformations and Imperforate anus/anal atresia in the immediate post-natal/neonatal period, followed by 2) long term medical management of sequelae of the congenital malformations e.g. renal and vertebral anomalies.

Even though optimal surgical correction is carried out in early life, some patients will continue to be affected by their congenital malformations throughout life.

V – Spinal surgery for vertebral anomalies and scoliosis (underlying costovertebral anomalies) are common. Vertical expandable prosthetic titanium rib (VEPTR) for treatment of thoracic insufficiency syndrome (TIS). Prophylactic de-tethering of spinal cord to prevent irreversible deficits (in filum terminale lipoma +/- low lying spinal cord) or at least halt progression of deformities (though not routinely offered if patient has anal atresia with tethered spinal cord). Laminectomy for back pain/scoliosis. Spinal cord stimulator. Occipital stabilisation. Cervical laminectomy and resection of posterior elements followed by stabilisation/fusion/traction (for congenital C5-6 dislocation).

A – Anal atresia or imperforate anus – surgery in first few days of life. Sometimes several surgeries for full reconstruction of intestine and anal canal. Imperforate anus – either complete correction immediately post birth or colostomy formation followed by reanastomosis and “pull-through” surgery. Atresia – repair without colostomy in infancy.

C – Ventricular septal defect, atrial septal defect, Tetralogy of Fallot, transposition of great arteries, truncus arteriosus correction. Ranges from anatomical abnormalities that do not require surgery to necessitating several stages of challenging surgery.

TE – Tracheoesophageal fistula repair / oesophageal atresia usually repaired in first few days of life with primary anastomosis (extrapleural approach/thoracoscopic/thoracotomy) unless other factors (long gap, low general condition, other major abnormalities) make this impossible, where a staged procedure is carried out (cervical oesophagostomy, abdominal oesophagostomy, ligation of distal oesophagus with gastrostomy and feeding jejunostomy). TEF repair almost always precedes repair of congenital heart disease if both are present.

Oesophageal stenosis – dilatation of oesophagus via balloon.

Tracheal stenosis – slide tracheoplasty.

Laryngo-tracheoesophageal cleft – conservative management for type 0 and 1; primary closure using either endoscopy or external surgery for types 2–4 (and if conservative management fails in type 0 and 1).

GOR – Nissen Fundoplication.

Diaphragmatic hernia repair.

Tracheomalacia – thoracoscopic aortopexy.

R – Surgery is done mainly to prevent damage from kidney and urological problems including persistent cloaca abnormalities i.e. dilatation of stenotic bulbar urethra.

Genitourinary abnormalities – reconstructive surgery, treated in a staged manner; primary repair (rectovaginal fistula, hypospadias)

L – Plastic surgery for polydactyly, syndactyly, wrist radialisation, but above all pollicisation for the severe degrees of thumb hypoplasia. Bilateral trochanteric surgery – hip dysplasia secondary to scoliosis.

Type of anaesthesia

General anaesthesia is needed for VACTERL association children due to the complexity of surgery undertaken. No reports of contraindications to either TIVA or volatile anaesthesia including nitrous oxide.

No reports of regional blocks being contraindicated in VACTERL patients.

Care must be taken when doing caudals or neuraxial blocks in VACTERL patients especially if they have an imperforate anus, genitourinary abnormalities or sacral dimple, as they frequently have occult spinal dysraphisms, e.g. tethered spinal cord, meningocele or lipo-myelomeningocele. When planning a neuraxial technique, a spine ultrasound should be performed. Difficult neuraxial block placement and failure has a higher incidence in patients with vertebral column abnormalities versus patients without spinal abnormalities. The utilisation of ultrasound has aided in accurate needle placement for neuraxial techniques when spinal pathology is present.

For thoracoscopic procedures, ventilation will need to be adjusted to normalise or allow for permissive hypercapnoea secondary to CO₂ insufflation. Also regular desufflation of thoracic cavity needs to be undertaken to restore sufficient oxygenation. Some patients with congenital heart disease may not tolerate the physiologic changes that accompany insufflation with thoracoscopic or laparoscopic procedures.

Necessary additional diagnostic procedures (preoperative)

Because of multiple systems affected by this association, baseline investigations are necessary to rule out or determine the severity of the condition in addition to the usual physical observations including height and weight.

Vertebral anomalies – X-ray, Ultrasound Scan (USS) and/or CT/MRI of the spine.

Anal Atresia – Physical examination/observation, abdominal ultrasound +/- additional testing for genitourinary anomalies (urethrocystography, urethroscopy, intestinal barium examination, abdominal/perineal USS, AXR).

Cardiac malformations – ECG (arrhythmias), echocardiogram +/- cardiac CT/MRI/angiogram (exclude cardiac or vascular abnormalities; evaluate structure and function of heart). Paediatric cardiology consultation. CXR (cardiomegaly).

Tracheoesophageal fistula – Physical examination/observation (contrast studies are rarely required); CXR – PA/lateral; CT/MRI (tracheoesophageal cleft) + CXR (aspiration evidence) and endoscopy; AXR (diaphragmatic hernia).

Renal anomalies – Renal ultrasound +/- voiding cystourethrogram; CT urogram multislice; abdominal CT; urine M/C/S.

Limb anomalies – Physical examination, X-rays (skeletal survey – limbs, hips, rib); angiography (radial artery hypoplasia).

Blood work – Full blood count, haemoglobin, haemocrit, electrolytes, renal function, liver and bone profiles, vitamin D, glucose, group and save, cross match; coagulation/bleeding time; chromosome analysis.

Post dialysis – blood urea, electrolytes, weight, (Pt with abnormalities of the heart and kidney may be more at risk for Haemolytic uremic syndrome).

Pulmonary workup – CXR +/- respiratory function tests.

Particular preparation for airway management

Airways can be anatomically challenging in VACTERL patients secondary to their craniofacial-vertebral deformities. A difficult airway trolley should be available in the operating room when anaesthetising these patients.

VACTERL children may have other associated abnormalities e.g. cleft lip and palate, hemifacial microsomia (hypoplastic mandible +/- webbed neck), and present for various types of surgery. A difficult airway and hence intubation should be anticipated.

Additionally extra equipment maybe needed for patients with cervical instability.

Extubating patients with difficult airways need to be taken with considerable caution. Depending on the surgery undertaken, some of these children will need postoperative intensive care ventilation for a few days, whilst others may be deemed safe to extubate on the operating table. History of an increased risk of aspiration, e.g. cleft lip/palate, tracheoesophageal fistula, may preclude early extubation.

There are reports of difficult intubation and tracheal damage in patients with oesophageal atresia. In patients with a TEF, endotracheal tube placement and mechanical ventilation can be especially challenging. This can be secondary to the proximity of the fistula to the carina, causing gastric insufflation, aspiration of gastric contents from the fistula, and prematurity, with an incidence of 30% in patients with a TEF. To properly position the ETT, an intentional mainstem intubation can be performed with subsequent withdrawal of the ETT just above the carina. Therefore, the ETT is positioned just above the carina and below the fistula, avoiding gastric insufflation and inadequate ventilation. This may not be possible in case of very large or pericarinal fistulas. Often, prior to intubation, rigid bronchoscopy is performed to define the anatomy, size and location of the fistula.

Isolated tracheal stenosis/atresia may exist which will cause impossible ETT intubation. Urgent tracheostomy will be needed. If laryngeal stenosis only is present, then smaller ETT

should be used rather than the correct size. Be careful in recently repaired laryngeal cleft when inserting ETT as it could damage the repair/reconstruction.

Particular preparation for transfusion or administration of blood products

Since there are major operations that will be undertaken in VACTERL infants, e.g. cardiac, TOF/OA, spinal operations, gastrointestinal and diaphragmatic hernia etc. – blood products need to be ordered and issued before the procedure is underway.

Particular preparation for anticoagulation

Patients may have coagulation issues secondary to chronic renal failure and being on dialysis. They may also be at a higher risk of developing uraemic haemolytic anaemia.

Otherwise, there are no reports on coagulation issues in VACTERL patients.

No particular reports for any anticoagulant regimen.

Particular precautions for positioning, transport or mobilisation

Not reported.

Probable interaction between anaesthetic agents and patient's long-term medication

Not reported.

Anaesthesiologic procedure

Avoid suxamethonium in patients with chronic renal failure, due to hyperkalaemic arrhythmias and arrest. Dialysis should be used to lower the potassium level prior to surgery.

No reports of either intravenous or inhalation anaesthetic agents being contraindicated.

Intravenous access can be challenging in patients with limb abnormalities.

For patients with TEF coming for repair, a rigid bronchoscopy is helpful to aid in ETT positioning. Anaesthesia with spontaneous ventilation versus positive pressure ventilation can avoid gastric insufflation through the fistula and subsequent pneumoperitoneum.

Patients with oesophageal atresia having had oesophageal reconstruction repair with a piece of colon, are at risk of aspiration including silent aspiration (up to 50%), as there is no lower oesophageal sphincter (LOS) present and motility is slow. Consider antacids +/- prokinetic agents and nil orally for 12 hours preoperatively. Rapid sequence induction (RSI) with adequate pre-oxygenation with elevation of head of the bed at induction of anaesthesia. Use clear mask so visualisation of aspiration is possible and have a large bore suction available.

Particular or additional monitoring

Forced air warming device should be used to prevent drop in the infant/neonate's (especially with low birth weight) core temperature.

Arterial and Central Venous Lines (CVL) are usually inserted in patients having prolonged operations, operations involving haemodynamic instability or major fluid shifts, e.g. cardiac and TOF/OA operations. Radial arterial line placement may be difficult or impossible if radial or upper limb anomalies are present.

NIRS (near infrared spectroscopy) can be used for cardiac and thoracoscopic repair of TOF/OA to ensure adequate oxygenation of cerebral tissues (although there are no consistent reports regarding this).

Possible complications

Patients with TOF/OA have a susceptibility to respiratory infections (atelectasis and pneumonia) due to a weakness in tracheal muscles and hyperreactive airways. These patients need active respiratory care with physiotherapy and antibiotics pre- and post-operatively.

Acute life threatening events (10–20%) may occur postoperatively in OA patients which include gastroesophageal reflux and tracheomalacia. In one study GOR occurred in 52% of the patients.

Patients that are premature (<1,500g) or have associated major cardiac conditions have an increased mortality rate post TOF/OA repair.

Patients with laryngeal clefts may present with chronic cough, aspirations, recurrent pneumonias and respiratory distress.

Postoperative care

Postoperative monitoring depends on type of surgery undertaken and preoperative medical condition.

Major operations, i.e. TOF/OA repair, laryngeal cleft, cardiac surgery, will require intensive care admission postoperatively for continued ventilation +/- paralysis, haemodynamic monitoring and appropriate fluid management, close observation for the development of potential complications as a result of surgery, feeding including total parental nutrition, antibiotics and analgesia.

Information about emergency-like situations / Differential diagnostics

Tracheal agenesis/stenosis with tracheoesophageal fistula can be associated with VACTERL association. Respiratory insufficiency in the newborn infant may be overcome by intubating the oesophagus or performing emergency tracheotomy and applying positive pressure ventilation. Success is not always possible.

Ambulatory anaesthesia

Not reported.

Obstetrical anaesthesia

Physiology during pregnancy can affect and even worsen the preexisting conditions of VACTERL parturients, mainly cardiac, spinal and respiratory/airway.

General, neuraxial (epidural +/- spinal) or a combined anaesthetic technique is possible after evaluating the issues pertinent to the individual patient. A full assessment needs to be undertaken in order to choose the optimal anaesthetic technique.

Beware that parturients may have a difficult airway and/or respiratory insufficiency (restrictive lung disease) due to major thoracocervical spinal abnormalities. The severity of the restrictive lung disease can negatively impact on ventilation +/- oxygenation during neuraxial or general anaesthesia. Airway assessment is important including Mallampati score. CXR and respiratory function tests will help to further define the extent of spinal abnormalities and any lung abnormalities, e.g. collapse, degree of lung expansion, respiratory reserve etc. Respiratory follow-up throughout pregnancy is important. Awake fibre optic intubation should be considered as a backup if direct laryngoscopy is felt to be dangerous.

Before a regional technique is undertaken, one needs to elicit with MRI where the spinal cord medulla ends and also define what spinal malformations if any, exist, e.g. segmentation in the lumbar region. Due to fixed lumbar scoliosis, it may be difficult to find lumbar landmarks for needle placement. Ultrasound can be used as an adjunct in this case (and it may reduce the incidence of dural puncture). Neuraxial anaesthesia can be used for both peri- and postoperative analgesia, although inadequate dermatomal block is a known complication.

ECHO is needed to rule out any abnormal cardiac morphology present. Cardiology follow-up for optimisation is important.

Literature and internet links

1. Al-Rawi O, et al. Oesophageal Atresia And Tracheo-Oesophageal Fistula; Continuing Education in Anaesthesia. Crit Care Pain 2007;7:1
2. Okada A, et al. Esophageal atresia in Osaka: A review of 39 years' experience. J Pediatr Surg 1997;32;11:1570–1574
3. Broemling N, Campbell F. Anesthetic management of congenital tracheoesophageal fistula. Paediatr Anesth 2011;21:1092–1099
4. Carli D, et al. VACTERL (vertebral defects, anal atresia, tracheoesophageal fistula with esophageal atresia, cardiac defects, renal and limb anomalies) association: disease spectrum in 25 patients ascertained for their upper limb involvement. J Pediatr 2014;164:458–462
5. Castori M, et al. Sirenomelia and VACTERL association in the Offspring of a Woman with Diabetes. Am J Med Genet 2010;152A:1803–1807
6. Castori M, et al. VACTERL association and Maternal Diabetes: a possible causal relationship? Birth Defects Res Part A-Clin Mol Teratol 2008;82:169–72
7. Camacho AF, et al. Monozygotic twins discordant for VACTERL association. Prenat Diagn 2008;28:366–368
8. Cote JC. A Practice of Anesthesia for Infants and Children. (Phil. : Els 2013) Philadelphia, PA (USA) Elsevier Publishing: 2013
9. van der Zee DC, et al. Thoracoscopic treatment of esophageal atresia with distal fistula and of tracheomalacia. Semin Pediatr Surg 2007;16:224–230
10. Jain D, Sampley S, Kaur G. Association of difficult airway to VACTERL anomaly: An anesthetic challenge. Anaesth pain intensive care 2013;17(2):192–194
11. Fargen KM, et al. Occipitocervicthoracic stabilization in pediatric patients. J Neurosurg 2011;8:57–62
12. Gedikbasi A, et al. Prenatal Diagnosis of VACTERL syndrome and Partial Caudal Regression Syndrome: a previously unreported association. J Clin Ultrasound 2009;37:464–466
13. Golonka NR, et al. Routine MRI evaluation of low imperforate anus reveals unexpected high incidence of tethered spinal cord. J Pediatr Surg 2002;37:966–969
14. Hilton G, et al. Anesthetic Management of a Parturient with VACTERL Association undergoing Cesarean delivery. Can J Anaesth 2013;60:570–576
15. Seo J, Kim DY, Kim AR et al. An 18-year experience of tracheoesophageal fistula and esophageal atresia. Korean J Pediatr 2010;53:705–710
16. Khatavkar SS et al. Anaesthetic Management for Cataract Surgery in VACTERL Syndrome. Case Report. Indian J Anaesth 2009;53:94–97
17. Krause U, et al. Isolated congenital tracheal stenosis in preterm newborn. Eur J Pediatr 2011;170:1217–1221
18. Kuo MF, et al. Tethered spinal cord and VACTERL association. J Neurosurg 2007;106(3rd Suppl):201–204
19. Leboulanger N, et al. Laryngo-tracheo-oesophageal clefts. Orphanet J Rare Dis 2011;6:81
20. Luce V, et al. Anesthetic management for a Parturient Affected by VACTERL Association. Anesth Analg 2004;98:874
21. Mariano ER, et al. Successful thoracoscopic repair of esophageal atresia with tracheaeso-phageal fistula in a newborn with single ventricle physiology. Anesth Analg 2005;101:1000–1002
22. O'Neill BR, et al. Prevalence of Tethered Spinal Cord in Infants with VACTERL. J Neurosurg Pediatr 2010;6:177–182
23. Pandey DB, et al. VACTERL Association. J Assoc Physicians India 2011;59:447–449
24. Raam MS, et al. Long-term outcomes of adults with features of VACTERL association. Eur J Med Genet 2011;54(1):34–41
25. Saade E, Setzer N. Anesthetic management of tracheoesophageal fistula repair in a newborn with hypoplastic left heart syndrome. Paediatr Anesth 2006;6:588–590
26. Schiffmann JH, et al. Tracheal agenesis, a rare cause of respiratory insufficiency in newborn infants. Monatsschr Kinderheilkd 1991;139:102–104
27. Solomon BD. VACTERL/VATER Association. Orphanet J Rare Dis 2011;6:56
28. Tandon RK, et al. Esophageal Atresia: Factors influencing survival – Experience at an Indian Tertiary Centre. J India Assoc Pediatr Surg 2008;13:2–6
29. Thaper A, et al. Bow-shaped tracheal rings: the lesson learnt from an endotracheal intubation. J Laryngol Otol 2004;118:732–733.

Last date of modification: October 2014

These guidelines have been prepared by:

Author

Elizabeth Richards, Anaesthesiologist, Kantonsspital Frauenfeld, Switzerland
lizzarichards@yahoo.com

Peer revision 1

Jennifer Dillow, Anaesthesiologist, University of New Mexico, Albuquerque, USA
jdillow@salud.unm.edu

Peer revision 2

Antonio Percesepe, Division of Medical Genetics, University Hospital of Modena, Italy
antonio.percesepe@unimore.it

Herausgeber



DGAI

Deutsche Gesellschaft
für Anästhesiologie und
Intensivmedizin e.V.
Präsident: Prof. Dr.
R. Rossaint, Aachen



BDA

Berufsverband Deutscher
Anästhesisten e.V.
Präsident: Prof. Dr.
G. Geldner, Ludwigsburg



DAF

Deutsche Akademie
für Anästhesiologische
Fortbildung e.V.
Präsident: Prof. Dr.
F. Wappler, Köln

Schriftleitung

Präsident/in der Herausgeberverbände
Gesamtschriftleiter/Editor-in-Chief:
Prof. Dr. Dr. Kai Zacharowski, Frankfurt
Stellvertretender Gesamtschriftleiter/
Deputy Editor:
Prof. Dr. T. Volk, Homburg/Saar
CME-Schriftleiter/CME-Editor:
Prof. Dr. W. Zink, Ludwigshafen

Redaktionskomitee/Editorial Board

Prof. Dr. G. Beck, Wiesbaden
Dr. iur. E. Biermann, Nürnberg
Prof. Dr. H. Bürkle, Freiburg
Prof. Dr. B. Ellger, Dortmund
Prof. Dr. K. Engelhard, Mainz
Prof. Dr. M. Fischer, Göttingen
Priv.-Doz. Dr. T. Iber, Baden-Baden
Prof. Dr. U. X. Kaisers, Ulm
Prof. Dr. W. Meißner, Jena
Prof. Dr. C. Nau, Lübeck
Dr. M. Rähmer, Mainz
Prof. Dr. A. Schleppers, Nürnberg
Prof. Dr. G. Theilmeier, Hannover
Prof. Dr. M. Thiel, Mannheim
Prof. Dr. F. Wappler, Köln
Prof. Dr. M. Weigand, Heidelberg

Redaktion/Editorial Staff

Carolin Sofia Kopp B.A. &
Dipl.-Sozw. Holger Sorgatz
Korrespondenzadresse: Roritzerstraße 27 |
90419 Nürnberg | Deutschland
Tel.: 0911 9337812 | Fax: 0911 3938195
E-Mail: anaesth.intensivmed@dgai-ev.de

Verlag & Druckerei

Aktiv Druck & Verlag GmbH

An der Lohwiese 36 |
97500 Ebelsbach | Deutschland
www.aktiv-druck.de

Geschäftsführung

Wolfgang Schröder | Jan Schröder |
Nadja Schwarz
Tel.: 09522 943560 | Fax: 09522 943567
E-Mail: info@aktiv-druck.de

Anzeigen | Vertrieb

Pia Engelhardt
Tel.: 09522 943570 | Fax: 09522 943577
E-Mail: anzeigen@aktiv-druck.de

Verlagsrepräsentanz

Jürgen Distler
Roritzerstraße 27, 90419 Nürnberg
Tel.: 0171 9432534 | Fax: 0911 3938195
E-Mail: jdistler@bda-ev.de

Herstellung | Gestaltung

Manfred Wuttke | Stefanie Triebert
Tel.: 09522 943571 | Fax: 09522 943577
E-Mail: ai@aktiv-druck.de

Titelbild

Dipl.-Designerin Monique Minde,
Nürnberg

Erscheinungsweise 2019

Der 60. Jahrgang erscheint jeweils zum
Monatsanfang, Heft 7/8 als Doppelausgabe.

Bezugspreise (inkl. Versandkosten):

• Einzelhefte	30,- €
• Jahresabonnement:	
Europa (ohne Schweiz)	258,- €
(inkl. 7 % MwSt.)	
Schweiz	266,- €
Rest der Welt	241,- €

Mitarbeiter aus Pflege, Labor, Studenten und Auszubildende (bei Vorlage eines entsprechenden Nachweises)

Europa (ohne Schweiz)	94,- €
(inkl. 7 % MwSt.)	
Schweiz	90,- €
Rest der Welt	94,- €

**Für Mitglieder der DGAI und/oder
des BDA ist der Bezug der Zeitschrift
im Mitgliedsbeitrag enthalten.**

Allgemeine Geschäfts- und Liefer- bedingungen

Die allgemeinen Geschäfts- und Liefer-
bedingungen entnehmen Sie bitte dem
Impressum auf www.ai-online.info

Indexed in **Current Contents®/Clinical
Medicine, EMBASE/Excerpta Medica;
Medical Documentation Service;
Research Alert; Sci Search; SUBIS
Current Awareness in Biomedicine;
VINITI: Russian Academy of Science.**

Nachdruck | Urheberrecht

Die veröffentlichten Beiträge sind urhe-
berrechtlich geschützt. Jegliche Art von
Vervielfältigungen – sei es auf mechani-
schem, digitalem oder sonst möglichem
Wege – bleibt vorbehalten. Die Aktiv
Druck & Verlags GmbH ist allein auto-
risiert, Rechte zu vergeben und Sonder-
drucke für gewerbliche Zwecke, gleich
in welcher Sprache, herzustellen. An-
fragen hierzu sind nur an den Verlag zu
richten. Jede im Bereich eines gewerbli-
chen Unternehmens zulässig hergestellte
oder benutzte Kopie dient gewerblichen
Zwecken gem. § 54 (2) UrhG. Die Wie-
dergabe von Gebrauchsnamen, Handels-
namen, Warenbezeichnungen usw. in
dieser Zeitschrift berechtigt auch ohne
besondere Kennzeichnung nicht zu der
Annahme, dass solche Namen im Sinne
der Warenzeichen- und Markenschutz-
Gesetzgebung als frei zu betrachten wä-
ren und daher von jedermann benutzt
werden dürften.

Wichtiger Hinweis

Für Angaben über Dosierungsanwei-
sungen und Applikationsformen kann
vom Verlag und den Herausgebern keine
Gewähr übernommen werden. Derartige
Angaben müssen vom jeweiligen An-
wender im Einzelfall anhand anderer
Literaturstellen auf ihre Richtigkeit über-
prüft werden. Gleiches gilt für berufs-
und verbandspolitische Stellungnahmen
und Empfehlungen.

Die Beiträge aus der A&I finden Sie online unter: www.ai-online.info

CONTACT US

Please do not hesitate to contact us. We will be glad to answer and provide further information to you at any time.

.....
Name

.....
First Name

.....
Department / Hospital

.....
Place

.....
Telephone

.....
E-Mail

.....
Date / Signature

Please contact me for further information

☐

I would like to participate in the project

☐

ADDRESS

German Society of Anaesthesiology and
Intensive Care Medicine
Ursula Homberg
Roritzerstrasse 27 | 90419 Nuremberg | Germany
Tel.: +49-911-9337828 | Fax: +49-911-3938195
Email: uhomberg@orphananesthesia.eu